

Index

- Abakaliki, mining contamination, 306
- Abidjan, 335
- Acid rain, 11
- Acidic deposition, 90
- Adsorption process, 183-185
- Aerosol concentrations in marine regions, 151
- Aerosol particles, 72, 149, 151, 168
dry deposition of, 160-163
- Aerosols, 45
atmospheric transport of, 107
solubility, 82
- Agricultural wastes, 226
- Air pollution, 336-337
- Air-to-sea exchange rates, 163-165
- Air-to-sea fluxes, 149, 166
- Algal growth, 228
- Algal products, 228
- Alkali industry, 327
- Alkylmercury fungicides, 60
- Alloy manufacturing, 77
- Alocasia odora*, 223
- Aminolevulinic acid dehydratase (ALA-D), 57
- Anaerobic conditions, 260
- Analytical data, quality control, 49
- Animal manure, 227, 228
- Antimony
deposition figures for, 110
in soils, 111
- Aquatic biota, 205-206
mercury bioaccumulation in, 255-277
- Aquatic ecosystems
Australian, 191, 202-207
management of, 206-207
- Aquatic environment
heavy metals in, 239
mercury in, rehabilitation and control measures, 269-272
trace metals in, 237-240
- Aquatic fauna, 205-206
- Arsenic
airborne, 63
alkyl species, 183
analytical concerns, 47
anthropogenic emissions, 294-297
anthropogenic sources, xix, 76-79, 84, 281-288
atmospheric, 74-76
atmospheric fluxes, 77
atmospheric transport, 81-83
behaviour in the body, 55-56
biogeochemical and biochemical pathways of, 44-45
biological methylation of, 44
chemical speciation from high-temperature processes, 79-81
chemicals, 286, 297
compounds, 47, 55
data needs, 45
emission rates from natural sources, 47
emissions, xxii, xxiii, 43, 44, 71-74, 280
equilibrium speciation calculations, 176
exposure thresholds, 64-65
general recommendations, 47
global atmospheric burden, 290-291
global atmospheric fluxes, 280, 291
global cycling of, 279-301
global oceanic burden and flux, 291-293
global tropospheric burden, 290
group report, 43-48
health effects and populations affected, xxiii, 46-47, 61-63
human exposure, 327-329
in aerosols, 45
in aquatic environment, 193-194
in coals and crude oils, 72
in drinking water, 46, 47, 62, 65, 328

Arsenic (*cont.*)

- in environment, 43, 317–320
- in fish, 46
- in foods, 63
- in India, 315–333
- in marine atmosphere, 162
- in marine rain, 152
- in medications, 63
- in oxygenated waters, 178
- in plants, 116
- in remote marine air and rain, 148
- in seafood, 226
- in sediments, 46, 294
- in soils, 45, 111, 289, 297, 319
- in vegetables, 328
- in water, 319
- industrial applications, 77
- ingestion, xxii, 43, 328
- inhalation, 43
- inorganic, 80
- long-term exposure to, 61
- low temperature volatilization, 289, 297
- major sources of, xix
- major transport pathway, 43
- metabolic factors associated with environmental exposure, 54
- microbial methylation, xx
- natural emissions, 297
- natural sources, 44, 78, 288–289
- net atmospheric flux, 291
- net oceanic flux, 294
- oceanic influx, 293
- organic, 80
- organs affected by, 62
- overt toxicity, 53
- pathways to man, 45
- pesticides, 77
- poisoning, xxii, 54
- residence times in seawater, 185
- route of entry, 43
- sinks for, 46
- solubility, 83
- sources, 43–44
- speciation and cycling in natural waters, 175–186
- speciation calculations, 179
- toxicity of, 63
- transfer rates in ecosystems, 45
- volatility of, 80

- volcanic source, 44

- Arsenic trioxide, 286

- Arsenious oxide, 319

- Aswan High Dam Reservoir

- khors, 236

- morphology, 236

- processes affecting distribution of trace metals in, 241

- residence time partitioning between particles and water, 241

- shape, 236

- siltation, 237, 247

- suspended matter transportation and movement patterns, 241

- thermal stratification, 242–243

- trace metals in, 235–254

- Atmospheric concentrations in

- Hawaii, 153–154

- Atmospheric emissions, xx

- Atmospheric transport, 81–83, 107–117

- Australia, heavy metal pollution, 188–216

- Background concentrations in air, rain and dry deposition, 149–152

- Backwaters, 326

- Bangalore, 326

- Biogeochemical cycling, 169, 170

- Biomass combustion, 13

- Blackfoot disease, 62, 63

- Blood samples, lead in, 224

- Bombay, 326, 327

- Brassica alboglabra*, 223

- Brassica chinensis*, 222, 227

- Brassica parachinensis*, 222, 223, 224, 227, 228

- Bubble Interfacial Microlayer Sampler, 164

- Cadmium

- airborne, 40

- and cigarette smoking, 55

- and zinc production, 38

- anthropogenic sources, 37–39, 76–79, 84

- atmospheric, 74–76

- in Samoa, 155

- atmospheric deposition, 128

- atmospheric fluxes, 77

- atmospheric transport, 81–83
- average daily intakes, 40
- behaviour in the body, 55–56
- biogeochemical cycling, 41
- body burden of, 40
- chemical forms, 80
- chemical speciation from high-temperature processes, 79–81
 - contamination, 38, 61
 - in elderly females, 61
- daily intake of, 328
- deposition in Norway, 108
- dietary intake, xxii, 39
- DTPA-extractable, 321
- emission control, 127
- emissions, 36, 37, 38, 71–74
- environmental dissipation, 40
- equilibrium speciation calculations, 176
- exposure of elderly females to, xix
- exposure thresholds, 64–65
- extraction of, 38
- from natural sources, 78
- gastrointestinal absorption, 55
- GEMS programme, 49
- geochemical behaviour of, 309
- geochemical dispersion, 304
- global cycling, 35
- group report, 35–42
- health effects and populations
 - affected, xxii, 35, 60–61
- human exposure, 327–329
- human exposure pathways, 39–40
- in Abakaliki area, 306
- in agricultural soils in taxonomic order, 124
- in air, 128, 323
- in aquatic environment, 193–194
- in Aswan High Dam Reservoir, 243
- in cigarette smoke, 40
- in coals and crude oils, 72
- in commercial crops, 127
- in common rocks, 122
- in crops, 41, 119–146, 128–140
 - from locations throughout US, 136–138
 - from major producing regions in US, 139
- in domestic water supplies, 121
- in environment, 320–324
 - in food chain, 120
 - in foodstuffs, 328
 - in India, 315–333
 - in kidney, 60
 - in lithosphere, 121
 - in marine atmosphere, 162
 - in marine sediments, 122
 - in natural surface soils, 114
 - in oxygenated waters, 178
 - in plants, 116
 - in remote marine air and rain, 148
 - in river waters, 246–247
 - in seafood, 226
 - in sediments, 122, 303–313
 - in sewage sludges, 127
 - in soils, 41, 111, 119–146, 303–313, 321
 - in surface soils, 123
 - in tropical soils, 307, 311
 - in vegetable crops, 132–134
 - in vegetative and reproductive tissues, 139
 - in water, 320
- inadvertent sources, 37
- industrial applications, 77
- information needs, 40–41
- ingestion of, 339
- inhalation, 121
- inventories, xxii
- levels near mining and smelting operations, 126–127
- metabolic factors, associated with environmental exposure, 54
- morality studies, 61
- natural levels in soils, 122–126
- natural sources, 37, 121–122
- overt toxicity, 53
- persistence, 120
- poisoning, xxii, 54, 120
- potential environmental hazards, 141
- principal uses, 39
- range of mean concentration by crop class, 140
- redistribution in soils, 311
- residence times in seawater, 185
- retention by organic matter, 124
- sensitivity to freshwater animals, 199
- solubility, 83

- Cadmium (*cont.*)
 sorption of, 322
 source inventories, 36–37
 sources and extent of
 contamination of soils,
 126–128
 speciation and cycling in natural
 waters, 175–186
 toxicity, 120
 vapour phase, 166
 volatility of, 80
 volcanic action, 37
 wastes, 39
- Calcutta, 326
- Cancer, 62
- Carbonate, 180
- Cement production, 77
- Chalcopyrite, 306
- Chandigarh, 320, 327
- Chembur, 326
- Chemical industries, 316–317
- Chicken manure, 226, 227, 229, 230
- Chloralkali industry, 327
- Chloralkali wastes, 93–97
- Chlorella pyrenoidosa*, 228
- Chlorella salina*, 228
- Chromium
 deposition patterns in Norway, 111
 in Lake Nubia sediments, 245
 in river sediments, 248
 in soils, 111
- Churchill River, 266, 267, 269
- Cigarette smoking, 40, 55, 121, 328
- Cladophora glomerata*, 10
- Coal combustion, 70–74, 80, 81,
 91–93, 284–285, 295, 297, 319
- Coal fired power plants, 127
- Coal production, 316–317
- Coastal waters, 319
- Cobalt
 in Aswan High Dam Reservoir,
 243
 in Lake Nubia sediments, 245
 in river sediments, 248
 in soils, 111
- Coke production, 76
- Combustion temperature, xx
- Cookson Reservoir, 266
- Copper
 deposition patterns in Norway, 111
 dry deposition of, 165
 in Aswan High Dam Reservoir,
 243
 in brown earth profiles, 113
 in dust, 223
 in fish, 230
 in natural surface soils, 114
 in river sediments, 248
 in river waters, 246–247
 in soils, 111
 mining, 201
 primary production of, 38–39
 refining, 281
 sensitivity to freshwater animals,
 199
 smelting, 281, 294, 295, 296,
 319
- Cows milk, 323
- Crassostrea gigas*, 226
- Cristigera spp.*, 10
- Cynodon dactylon*, 223, 225
- Cyprinus carpio*, 229
- Daphnia magna*, 10
- Desiccants, 286
- Developing countries, 13–14,
 335–341
 cadmium contamination, 141
 mercury in, 21
 monitoring heavy metals, 343–347
- Developing countries, xix
- Drinking water, 46, 47, 63, 319, 328
 arsenic in, 62, 65
- Dry deposition
 measurement of, 160–163
 model calculations of, 160
- Ecosystem studies, 199–202
- Egypt, *see* Aswan High Dam
 Reservoir
- Electroplating wastes, 188
- Electrostatic precipitators, 296
- Eleusine indica*, 223, 225
- Energy production, 296
- Environmental compartments, xxi
- Environmental contamination, 15
- Environmental cycling, 15
- Environmental degradation, 15
- Environmental fluxes, xxiii
- Environmental sinks, xxi
- Equilibrium speciation calculations,
 176

- Equilibrium speciation models, 176
Equisetum, 10
Erie, Lake, 326
Erythrocyte protoporphyrin
 fluorescence, 13
Estuarine waters, 319, 326

Fertilizers, 127-128, 318, 321,
 338
Finniss River, 197, 201
Fish
 arsenic in, 46
 heavy metals in, 250-251
 lead in, 230
 mercury in, 25-26, 49, 59, 90,
 257-268, 326
 methylmercury in, 25, 268
 production and contamination,
 229-230
 zinc in, 230
Fish studies, Nile River, 248-252
Flow variability, 202-203
Fly ash, 320, 323
Foninalis sp., 10
Food additives, 286
Food chain, xix, 3, 17, 18, 120, 121,
 128
Food contamination, 63
Foodstuffs, 39, 319, 320, 323, 328
Forest fires, 289
Fossil fuels, xx, 79, 80, 337
Fowler's solution, 62, 63
Free erythrocyte protoporphyrin
 (FEP), 57
Freshwater flora, 206
Freshwater invertebrates, 205
Fulvic acids, 180
Fungicides, 60, 319, 327

Galena, 306, 339
Gangrene, 62
Gasoline combustion, 79, 81,
 219-224, 336
GEMS programme, 49
Geothermal power, 296
Ghana, 339
Glass manufacture, 288
Global mass balance, 148
Goa, 319
Grain, mercury in, 326
Groundwaters, 338

Haemangioendothelioma of the liver,
 62
Hard and soft acid and base (HSAB)
 concept, 180
Harmattan season, 339
Hawaii, atmospheric concentrations
 in, 153-154
Hearing loss, 63
Heavy metals
 air pollution, 336-337
 analytical methods, 197
 anthropogenic sources, 165, 188,
 336
 biogeochemical cycling of, 188,
 191-197
 chemical analysis, 196-197
 chemical forms, 239
 ecological effects, 199
 effects on Australian ecosystems,
 188
 effects on biota, 198-202
 fate on entering the ocean,
 167-169
 in animals, 115-116
 in aquatic environment, 239
 in fish, 250-251
 in marine atmosphere, 166-167
 in marine biota, 192
 in marine organisms, 226
 in ombrotrophic bogs, 115
 in plants, 115-116
 in sediments and waters, 192
 in soils, 111-115, 308-309
 in surface waters, 337
 in terrestrial environment, 192
 levels of, 191-192
 long-range atmospheric transport,
 107-117
 monitoring, 335-341
 natural sources, 336, 338-339
 pollution in Australia, 188-216
 pollution sources, 336-339
 sources of, 188-190
 speciation studies, 196-197
 toxicity studies, 198-199
 transfer across air-sea interface,
 147-173
 transfer process, 195, 197-198
Herbicides, 286
High alkalinity river water (HARW),
 177, 180

- High temperature processes on
power generation and industry,
69–87
- Hong Kong
general background, 218–219
lead contamination, 217–233
traffic density, 219
- Human hair, lead in, 224
- Humic acids, 180
- Hydroelectric power plant, 266
- Hydroelectric reservoirs, 255–277
- Hydrolysis constants, 184
- Hydrolysis sequence, 180, 184
- Hylocomium splendens*, 10, 111
- Hyperkeratosis, 61
- Hyperpigmentation, 61
- Hypertension, 328
- Hypnum cupressiforme*, 10
- Ibadan, 337
- India
arsenic in environment, 317–320
cadmium in environment, 320–324
chemical industries, 316–317
cycling of arsenic, cadmium, lead
and mercury in, 315–333
human exposure to arsenic,
cadmium, lead and mercury,
327–329
lead in environment, 324–325
mercury in environment, 325–327
mineral processing, 316–317
perspectives, 329–340
- Indra Prastha power station, 319,
320
- Industrial development, 329
- Industrial emissions, 337
- Industrial processes, 69–87
- Industrial production, 317
- Industrial sources, 190, 225
- Industrial wastes, 7, 188, 190
- Industrialization, 335
- Iron
in sediments, 307–308
manufacturing, 76, 77, 80
- Iron ore processing, 295
- Iron ore tailings, 226
- Itai-itai disease, 61
- Ivory Coast, 335
- King River 201
- La Grande Reservoir, 265
- Lagos, 335, 336
- Lakes, 204–205
acidification, 90
quantitative effect of sediment, 243
trophic status, 27
- Land clearance, 286
- Lead
accumulation, xxi
adverse effects in children, 14
alkyl species, 183
and urban microenvironment,
12–13
anthropogenic, xix, xxi, 14, 76–79,
84
as function of particle size, 162
at open ocean sites, 167
atmospheric, 74–76, 167
in Samoa, 155
atmospheric fluxes, 77
atmospheric transport, 81–83
behaviour in the body, 55–56
biogeochemical cycle, 3, 4
biomagnification, 7
biomethylated, 9
biopurification 7
biotransformation of inorganic
environmental, 4
blood levels, 13, 224
brain damage, xxi
chemical speciation from high-
temperature processes, 79–81
concentration in ice layers at
North Pole, 3
cycling, 5
factors affecting, 10–11
daily intakes, 329
deposition in Norway, 108
dry deposition, 165
dust vector, 13
emission controls, 11
emissions, xxi, 8, 9, 13, 71–74,
189, 219–220, 336
environmental hazard, 3
environmental pathways, 5
equilibrium speciation calculations,
176
exposure thresholds, 64–65
fishing weights, 7
from gasoline combustion, 219–224
from industrial activities, 225

- from natural sources, 78
 - from sediments, 7
 - from water pipes, xix
 - gasoline additives, 8
 - GEMS programme, 49
 - geochemical behaviour, 309
 - geochemical dispersion, 304
 - group report, 3-16
 - haematological effects, 56
 - health effects and populations
 - affected, xix, 3, 56-59
 - human exposure, 327-329
 - in Abakaliki area, 306
 - in air, 5, 219-220, 323
 - in aquatic environment, 193-194
 - in brown earth profiles, 113
 - in central nervous system, 57
 - in children, 3, 12, 54, 339
 - in coals and crude oils, 72
 - in consumer goods, 14
 - in developing countries, 13-14
 - in domestic animals, 327
 - in dust, 220-223
 - in environment, 324-325
 - in fish, 230
 - in freshwater, 5, 10
 - in gasoline, 11, 14
 - in haematological system, 56
 - in Hong Kong, 217-233
 - in human hair, 224
 - in India, 315-333
 - in marine atmosphere, 162
 - in natural surface soils, 112, 114
 - in nervous system, 56
 - in ocean water, 7, 185
 - in oxygenated waters, 178
 - in paints, 225
 - in plants, 116, 223-224
 - in rain, 150
 - in remote marine air and rain, 148
 - in renal system, 56
 - in river sediments, 248
 - in river waters, 246-247
 - in salt water ecosystems, 6
 - in sediments, 225, 303-313
 - in soils, 220-222, 303-313, 321, 324
 - in surface peat, 114
 - in tropical soils, 307, 312
 - in vegetables, 227
 - in vegetation, 223
 - in water, 325
 - indicator organisms, 10
 - industrial applications, 77
 - ingestion of, 339
 - inhalation, 54
 - inorganic form, 4
 - major sources of, xix
 - mass-size function, 162
 - metabolic factors associated with environmental exposure, 54
 - monitoring, 12
 - national level data, 11
 - on land, 5
 - organometallic compounds, 4
 - overt toxicity, 53
 - oxide, 80
 - pathways and accumulating compartments, 9
 - poisoning, xxi, 55
 - primary production, 38-39
 - regional information, 7-8
 - regional inventories, 15
 - reservoirs of, 9
 - residence times in seawater, 185
 - responses of biota, 10
 - riverborne, 7
 - routes of human exposure, 5
 - shot ingestion, 7
 - smelting, 283-284
 - sources, xix, 8
 - speciation and cycling in natural waters, 4, 175-186
 - susceptibility of children to, 58
 - susceptibility of humans to, xxi, 14
 - technologies shift, xxii
 - terrestrial food chains, 3
 - transplacental transfer, 55
 - via wet, dry or cloud deposition, 5
 - volatility of, 80
 - water pipes, 13
- Lead chloride, 80
- Lignite deposits, 316
- Limnocostracans, 269
- Litter
 - accumulation, 338
 - decomposition, 10
- London, 336
- Low alkalinity river water (LARW), 177, 180
- Lung cancer, 62, 63

- Magela Creek, 197
- Manganese
- in Aswan High Dam Reservoir, 241, 243
 - in Lake Nubia sediments, 245
 - in river sediments, 248
 - in sediments, 307-308
- Manures, 318
- Marine aerosols, 169
- Marine atmosphere. heavy metals in, 166-167
- Marine environment, micro-organisms in, 166
- Mean oceanic residence time, 184
- Mercury
- adsorption, 92
 - airborne, 99
 - analytical sampling, 22
 - and environmental health, 19
 - and selenium relationship, 28-29
 - anthropogenic fluxes, 20
 - anthropogenic sources, 76-79, 84
 - aquatic environment, 17
 - aquatic food chain, 19
 - as fungicide, 60
 - atmospheric, 74-76, 83
 - atmospheric deposition rates, 97
 - atmospheric fluxes, 77
 - atmospheric load, 18
 - atmospheric transport, 81-83
 - bacterial methylation, 26
 - behaviour in the body, 55-56
 - bioaccumulation, 18
 - by aquatic biota, 255-277
 - bioamplification, 259
 - biogeochemical cycle, 20-21, 90
 - biological impacts, 19
 - biomagnification, 17, 19
 - bulk deposition, 95
 - chemical speciation from high-temperature processes, 79-81
 - concentrations in various atmospheres, 96
 - condensation, 92
 - current trends in use, 21
 - demethylation, 24, 25
 - deposition, 22, 26-27, 90-91
 - early toxic effects, 17
 - emissions, xx, 18, 20, 27-28, 71-74, 90-91
 - from coal-fired power plants, 93
 - from waste deposits, 96
 - equilibrium concentration in pure water, 102
 - equilibrium speciation calculations, 176
 - from natural sources, 78
 - from solid wastes, 93
 - gaseous, 81, 163, 167
 - general atmospheric cycle, 100
 - global atmospheric pool, 17
 - global cycling, 90, 99-102
 - global dispersion, 17
 - global wet deposition, 102
 - group report, 17-33
 - hair analysis, 19
 - health effects and populations affected, 18, 59-60
 - human exposure, 327-329
 - in air, 95, 100
 - in aquatic environment, 193-194
 - rehabilitation and control measures, 269-272
 - in atmosphere, 21, 22
 - in coals and crude oils, 72
 - in developing countries, 21
 - in domestic animals, 327
 - in environment, 23, 325-327
 - in fish, 25-26, 49, 59, 90, 257-268, 326
 - in grain, 326
 - in India, 315-333
 - in muscle, 25
 - in oxygenated waters, 178
 - in plants, 99
 - in sediments, 326
 - in soils, 97-99
 - in stored solid wastes, 27-28
 - in surface soils, 113
 - in water, 325
 - industrial applications, 77
 - inorganic compounds, 99
 - lake budget, 27
 - long range transport, 17
 - major pathway to man, 17
 - major sources of, xix
 - marine biological sources, 152
 - marine pollution, 21
 - metabolic factors, associated with environmental exposure, 54
 - methylation, xx, 17, 24, 25, 27
 - microbial demethylation, 24

- monitoring, 21
- natural emission rates from solid materials, 95
- organic compounds, 99
- overt toxicity, 53
- particulate, 95, 166
- plant uptake studies, 98-99
- poisoning, xx, 60, 339
- puzzles regarding, xxii
- removal from atmosphere, 101
- residence times in seawater, 185
- sampling, 99
- solubilities of, 102
- speciation and cycling in natural waters, 175-186
- technologies shift, xxii
- total annual surface emission, 96
- toxic effects, 19
- toxicity modification, 29
- vapour phase, xx, 22, 80, 89-106, 166
- volatility of, 80, 93
 - see also Methylmercury
- Mercury cycle, 21
- Mercury simicides, 21
- Metabolic factors, 54-56
- Metal alkyls, 180
- Metal-organic complexes, 196, 308
- Metal-organic interactions in natural waters, 181
- Metals Cycling Workshop, xx
- Metastable oxidation states, 180
- Meteorological factors, 148
- Methylmercury
 - anoxic systems, 183
 - exposure thresholds, 64-65
 - in fish, 25, 268
 - in surface water, 24
 - intoxication, 59
 - poisoning, 18, 29, 54
 - toxicity, 29
 - transplacental transfer, 55
 - uptake efficiency from water, 257
- Microbial redox processes, 181
- Micrococcus luteus*, 10
- Mineral deposits, 304
- Mineral processing, 316-317
- Mineral production, 318
- Mineral reserves, 318
- Mining operations, 126, 189, 197, 200, 201, 306, 337
- Molonglo River, 200
- Monitoring
 - comprehensive, 340
 - coordinating body, 346
 - environmental, 13, 339-340
 - funding limitations, 12
 - handbooks of agreed techniques, 345
 - heavy metals, 335-341, 343-347
 - historical record, 12
 - international collaboration and financial aid, 344
 - international compatibility, 344
 - international funding, 346
 - planning, 12
 - programme proposals, 344
 - specialist training for, 346
 - three-tier system, 344-345
- Mugil dues umierie*, 326
- Murray River, 203
- Mytilus edulis*, 10
- Nasal septum lesions, 61
- Nasser, Lake, 242-245
- Natural gas flaring, 337
- Natural sources, unknown, 165
- Natural waters, metal-organic interactions in, 181
- Nelson River, 266
- New York, 336
- Nickel
 - deposition patterns in Norway, 111
 - in soils, 111
- Nigeria, 335, 337, 339
 - dispersion of cadmium, lead and zinc in soils and sediments, 303-313
- Nile River
 - ecosystems, 235-254
 - fish studies, 248-252
 - monitoring sites, 240
 - pollution assessment, 248-252
 - trace metals in, 245-246, 248, 249
 - watershed areas, 237
- Nitrogen fertilizers, 127
- Non-equilibrium behaviour, 180
- Non-ferrous metal production, 74-76, 79, 80, 126, 140, 294, 296-297

- Norway
 deposition of heavy metals,
 107–117
 heavy metals in soils, 111–115
 sampling stations, 109
 trace element deposition patterns,
 109–111
- Nubia, Lake, 242, 243, 245
- Nutritional deficiencies, 14
- Nutritional factors, 19
- Oat grain, 130
- Ocean cycles, 148
- Oceans
 arsenic influx and outflux, 293–294
 heavy metals in, 167–169
- Oil combustion, 70–74, 80
- Ombrotrophic bogs, heavy metals in,
 115
- Opinacca reservoir, 265
- Opium, 328
- Organic complex formation, 180
- Organomercurial compounds, 19,
 327
- Oxygen redoxcline in stratified lakes,
 243
- Paints, lead-containing, 225
- Peat profiles, 115
- Petroleum products, 325
- Phosphorus fertilizers, 127–128
- Pig manure, 227, 229, 230
- Pinna bicolor*, 200
- Plants, lead in, 116, 223–224
- Pleurozium schreberi*, 10
- Port Harcourt, 337
- Power generation, 69–87
- Power plants, 337
- Precipitation scavenging, 101, 102
- Precipitation systems, 152
- Proteinuria, 60
- Punjab, 320, 323
- Quebec reservoirs, 265
- R-mode factor analysis, 308, 312
- Radionuclides, 168
- Rain collection, for metal analysis,
 152
- Rain concentrations, 149–150
 collected from islands and ships,
 150
 marine, 150
- Rain fluxes, and washout factors,
 156–159
- Rain samples
 collection from ships, 155
 collection system, 154
 contamination of, 154–156
 ocean surface, 154
 representativeness of, 152
- Raphanus sativus*, 227
- Recycled components for dry
 deposition measurements, 165
- Recycled metals in marine rains, 164
- Redox speciation, 180–183
- Reference standards, 50
- Refuse compost, 227
- Refuse incineration, 76, 77, 80
- Reservoirs, mercury in fish from,
 258–268
- Residence times, 183–185
- River sediments, trace metals in,
 248, 249
- River waters, trace metals in,
 246–247
- Rivers, 202, 203, 206, 207
- Saccostrea glomerata*, 226
- Saharan dust, 339
- Samoa, lead and cadmium in rain
 samples, 155
- Sarotherdon mossambicus*, 230
- Scavenging ratios, *see* Washout
 factors
- Scavenging residence times, 184
- Scrubbing agents, 239
- Sea salt particle deposition, 161
- Sea salt spray, 288, 294
- Sea spray recycling, 163
- Seawater, 167, 177, 179, 180, 325
 residence times in, 185
 speciation models, 178
- Selenium
 and mercury relationship, 28–29
 in plants, 116
 in surface soils, 113
- Sewage dumping, 225–226
- Sewage effluents, 190
- Sewage irrigation, 321, 322
- Sewage sludge, 41, 127, 133, 134,
 139, 190, 226–230, 323, 338
- Sewage system, 7, 39

- Skin tumours, 61
Smelting operations, 126, 281,
283–284, 294–296, 319, 337
Soils
 arsenic in, 45, 111, 289, 297, 319
 cadmium in, 41, 111, 119–146,
 303–313, 321
 chromium in, 111
 cobalt in, 111
 copper in, 111
 ferrallitic, 309
 heavy metals in, 111–115, 308–309
 lead in, 220–222, 307, 312, 321,
 324
 mercury in, 97–99, 113
 mercuriferous, 97–99
 nickel in, 111
 oxidation–reduction potential of,
 133
 pH effects, 128–134, 322
 pollution of, 338
 roadside, 336
 sewage-irrigated, 321, 322
 trace elements in, 306
 tubewell-irrigated, 322, 324
 waste water-irrigated, 324
 zinc in, 111, 303–313
South Esk River, 200–201
Speciation studies, 196–197
Spencer's Gulf, 200
Sphagnum moss, 269
Sphalerite, 306
Stability constants, 176–177
Standard reference material, 49–50
Steel manufacturing, 76, 77, 80, 284
Stormwater quality, 190
Stream sediments, 304, 307, 311, 312
Streams, 202, 203, 206, 207
Sulphate complexation, 179
Sulphide forming elements, 71
Sulphide ores, 74
Sulphur compounds, 260
Surface complexation constants, 184
Surface complexation model, 183
Swiss chard, 130
Synfuel technology, 296

Tilapia mozambique, 326
Tillandsia usneoides, 10
Tin mining, 200
Tobacco, 39

Total dissolved solids (TDS), 203,
204
Toxicity studies, 198–199
Trace metals, 71, 73, 77, 81, 82
 adsorbability and relative residence
 time, 239
 and suspended solids, 247–248
 atmospheric concentrations of, 148
 biochemical role, 239
 biogeochemical cycle, 239
 deposition patterns in Norway,
 109–111
 dry deposition of, 160
 in aquatic environment, 237–240
 in Aswan High Dam Reservoir,
 235–254
 in Aswan High Dam Reservoir
 sediments, 245–246
 in Nile River, 245–246, 248, 249
 in river sediments, 248, 249
 in river waters, 246–247
 in rock–soil–plant systems, 304
 in sediments, 309–312
 in surface waters, 169
 in tropical soils, 309–312
 ingestion, 239
 mass median diameter of, 82
 mass-size distribution of, 160
 non-biodegradable, 239
 point and non-point sources, 189
 regional distribution of, 304
 removal from atmosphere, 82
 sampling of precipitation for, 108
 suspended solids as scavengers for,
 247
Transfer processes, 197–198
Tropical ecosystems, 303–313
Tropical rain forests, 338
Tropical rivers, 203
Tropical weathering, 338

Ulva lactuca, 226, 228
United States, cadmium levels in
 soils and crops, 119–146
Uranium mining, 197, 201
Urban stormwater samples, 190
Urbanization, 335

Vanadium in seawater, 167
Vegetables
 arsenic in, 328

- Vegetables (*cont.*)
 cadmium in, 132–134
 lead in, 223, 227
Volcanoes, 289
- Washout factors, 166
 and mass median diameter
 (MMD), 157
 and rain fluxes, 156–159
 defined, 156
 to calculate metal deposition rates,
 156
- Waste disposal, 39
Waste incineration, 165, 288, 337
Waste materials, 93–97
 recycling of, 226–230
Water pollution, 325, 337–338
Water quality, 203–204
Water supplies, cadmium in,
 121
West Africa, 335–341
Willard Bay reservoir, 258
Wind blown dust, 189, 339
Wind erosion, 288
Wind flow patterns, 151
- Wolfram mining, 200
Wood fuel consumption, 286
Wood preservatives, 287, 297
- Zinc
 dry deposition, 165
 geochemical behaviour, 309
 geochemical dispersion, 304
 in Abakaliki area, 306
 in Aswan High Dam Reservoir,
 243
 in dust, 223
 in fish, 230
 in Lake Nubia sediments, 245
 in natural surface soils, 114
 in plants, 116
 in river sediments, 248
 in river waters, 246–247
 in sediments, 303–313
 in soils, 111, 303–313
 in tropical soils, 307
 sensitivity to freshwater animals,
 199
Zinc pollution, 200
Zinc production, 74–75, 284